

1

CAPTURE PURIFICATION PROCESSES FOR PROTEINS EXPRESSED IN A NON-MAMMALIAN SYSTEM

This application is a continuation of U.S. application Ser. No. 14/599,336, filed on Jan. 16, 2015, now U.S. Pat. No. 9,643,997; which is a divisional of U.S. application Ser. No. 12/822,990, filed on Jun. 24, 2010, now U.S. Pat. No. 8,940,878; which claims the benefit of U.S. Provisional Application No. 61/220,477 filed Jun. 25, 2009, which is incorporated by reference herein.

FIELD OF THE INVENTION

The present invention relates generally to processes for purifying proteins expressed in non-mammalian systems in both non-native soluble and non-native insoluble forms, and more particularly to the direct capture of such proteins from a refold mixture or a cell lysate pool by a separation matrix.

BACKGROUND OF THE INVENTION

Fc-containing proteins are typically expressed in mammalian cells, such as CHO cells. The use of affinity chromatography to purify Fc-containing proteins is documented (see, e.g., Shukla et al., (2007) *Journal of Chromatography B* 848(1):28-39) and is successful, in part, due to the degree of Fc structure observed in proteins expressed in such systems. Fc-containing proteins expressed in non-mammalian cells, however, are often deposited in the expressing cells in limited solubility forms, such as inclusion bodies, that require refolding, and this has been a limiting factor in selecting non-mammalian systems for expressing Fc-containing proteins.

A drawback to the use of Protein A, Protein G and other chemistries is that in order for a protein comprising an Fc region to associate with the Protein A or Protein G molecule, the protein needs to have a minimum amount of structure. Often, the requisite amount of structure is absent from proteins expressed recombinantly in a soluble, but non-native, form and consequently Protein A chromatography is not performed in a purification process.

In the case of a protein expressed in an insoluble non-native form, Protein A chromatography is typically not performed in a purification process until after the protein has been refolded to a degree that it can associate with the Protein A molecule and has been subsequently diluted out of its refold solution. This is because it was believed that after a protein has been refolded it was necessary to dilute or remove the components of the refold mixture in a wash step, due to the tendency of the components that typically make up a refold solution to disrupt interactions between the target protein and the Protein A molecules (Wang et al., (1997). *Biochem. J.* 325 (Part 3):707-710). This dilution step can consume time and resources which, when working at a manufacturing scale of thousands of liters of culture, can be costly.

The present disclosure addresses these issues by providing simplified methods of purifying proteins comprising Fc regions that are expressed in non-mammalian expression systems in a non-native soluble form or in a non-native insoluble form.

SUMMARY OF THE INVENTION

A method of purifying a protein expressed in a non-native soluble form in a non-mammalian expression system is

2

provided. In one embodiment the method comprises (a) lysing a non-mammalian cell in which the protein is expressed in a non-native soluble form to generate a cell lysate; (b) contacting the cell lysate with an separation matrix under conditions suitable for the protein to associate with the separation matrix; (c) washing the separation matrix; and (d) eluting the protein from the separation matrix.

The protein can be a complex protein, such as a protein is selected from the group consisting of a multimeric protein, an antibody and an Fc fusion protein. The non-mammalian expression system can comprise bacteria or yeast cells. The separation matrix can be an affinity resin, such as an affinity resin selected from the group consisting of Protein A, Protein G and a synthetic mimetic affinity resin, or it can be a non-affinity resin, such as a non-affinity resin selected from the group consisting of ion exchange, mixed mode, and a hydrophobic interaction resin. The cell lysate can be filtered before it is contacted with the separation matrix. Although not required, the method can further comprise refolding the protein to its native form after it is eluted from the separation matrix.

A method of purifying a protein expressed in a non-native limited solubility form in a non-mammalian expression system is provided. In one embodiment that method comprises (a) expressing a protein in a non-native limited solubility form in a non-mammalian cell; (b) lysing a non-mammalian cell; (c) solubilizing the expressed protein in a solubilization solution comprising one or more of the following: (i) a denaturant; (ii) a reductant; and (iii) a surfactant; (d) forming a refold solution comprising the solubilization solution and a refold buffer, the refold buffer comprising one or more of the following: (i) a denaturant; (ii) an aggregation suppressor; (iii) a protein stabilizer; and (iv) a redox component; (e) applying the refold solution to a separation matrix under conditions suitable for the protein to associate with the matrix; (f) washing the separation matrix; and (g) eluting the protein from the separation matrix.

The non-native limited solubility form can be a component of an inclusion body. The protein can be a complex protein, such as a complex protein selected from the group consisting of a multimeric protein, an antibody, a peptibody, and an Fc fusion protein. The non-mammalian expression system can be bacteria or yeast cells. The denaturant can comprise one or more of urea, guanidinium salts, dimethyl urea, methylurea and ethylurea, the reductant can comprise one or more of cysteine, DTT, beta-mercaptoethanol and glutathione, the surfactant can comprise one or more of sarcosyl and sodium dodecylsulfate, the aggregation suppressor can be selected from the group consisting of arginine, proline, polyethylene glycols, non-ionic surfactants, ionic surfactants, polyhydric alcohols, glycerol, sucrose, sorbitol, glucose, tris, sodium sulfate, potassium sulfate and osmolytes, the protein stabilizer can comprise one or more of arginine, proline, polyethylene glycols, non-ionic surfactants, ionic surfactants, polyhydric alcohols, glycerol, sucrose, sorbitol, glucose, tris, sodium sulfate, potassium sulfate and osmolytes, and the redox component can comprise one or more of glutathione-reduced, glutathione-oxidized, cysteine, cystine, cysteamine, cystamine and beta-mercaptoethanol. The separation matrix can be an affinity resin such as an affinity resin selected from the group consisting of Protein A, Protein G, and synthetic mimetic affinity resin or the separation matrix can be a non-affinity resin selected from the group consisting of ion exchange, mixed mode, and a hydrophobic interaction resin.